

Clark, A. J., Crawford, C. H., King, B. D., Boman, A. M., and Lyons, T. A. 2016. Material properties of hagfish skin, with insights into swimming behavior. <i>Biol Bull</i> 288, 241-256.	The authors study material properties of hagfish skin, develop stress-strain curves, and analyze how the results play a role in hagfish swimming abilities.	http://www.journals.uchicago.edu/doi/pdfplus/10.1086/6881.c230a2510				
Clark, A. J. and Summers, A. P. 2012. Ontogenetic scaling of the morphology and biomechanics of the feeding apparatus in the Pacific herring <i>Clupea pallasii</i> . <i>Journal of Fish Biology</i> 80, 36-59.	This paper presents measurements of mouth area, feeding apparatus length, basal plate size, theoretical dorsal plate remanile force, penetration force, and optimal tooth stress.	http://dx.doi.org/10.1111/j.1095-8649.2011.02412.x				
Fudge, B. A., Levin, N., Fink, A., and Grallion, J. M. 2003. Composition, morphology, and mechanics of Hagfish slime. <i>JEB</i> 208, 4613-4625.	This article looks at the composition of slime and presents mechanical properties of the slime's main component.	https://www.researchgate.net/profile/Scott_Chia/publication/744229_Composition_Morphology_and_Mechanics_of_Hagfish_Slime/files/5456d806f01b6c6e698e/Composition-Morphology-and-Mechanics-of-Hagfish-Slime.pdf				
Kier, W. M. 1992. Hyostatic skeletons and muscular hydrostoms. <i>BioScience</i> 41, 202-210.	This article discusses the nature of hyostatic skeletons.	http://leba.bio.umc.edu/kier/pdf/kier_1992.pdf				
Kier, W. M. and Smith, K. K. 1985. Torques, stresses and strains: the biomechanics of movement in muscle-hydrostatic. <i>Zoological Journal of the Linnean Society</i> 81, 291-324.	This article looks at the biomechanics of muscular hydrostoms.	http://leba.bio.umc.edu/kier/pdf/kier_1985.pdf				
Lim, J. L. 2013. Kinematics and hydrodynamics of undulatory locomotion in hagfishes (<i>Ophiodon elongatus</i>) and hagfish-like robotic models. <i>Harvard Undergraduate Journal of Science</i> 14, 149-169.	This dissertation examines the kinematics of hagfish swimming motion and compares it to model swimming performance using robot models.	https://dash.harvard.edu/bitstream/handle/1:111252201/lim_gsj_sci_0883.pdf?sequence=1				
Long, J. J., H. Koob-Franz, M., Sivewill, R., and Koob, T. J. 2007. The rheology of hagfish <i>Rosalia glaucosa</i> : tissue elastic properties and mechanical functions during steady swimming. <i>J Exp Biol</i> 280, 2193-2201.	This study looks at locomotor function of the hagfish notochord through the measurement of flexural stiffness and flexural damping under <i>in vivo</i> conditions.	http://jeb.biologists.org/content/280/24/2193.full				
Parrish, J. K., Kawan, S., Herberich, G., Doherty, J., and Strubbe, A. 2001. Localization of breakage points in knotted string. <i>New Journal of Physics</i> 3, 101-103.	This paper uses high speed cameras to identify the breakage point in knotted string under tension.	http://dx.doi.org/10.1088/1464-4201/3/1/010103				
Sambrook, C., Flaxell, G., Flaxell, T., and Hochberg, R. 2006. Octopuses use a human-like strategy to control precise point-to-point arm movements. <i>Curr Biol</i> 16, 387-392.	This paper studies the biomechanics of point-to-point movement in octopuses.	http://www.sciencedirect.com/science/article/pii/S096922600627247				
Turner, J. C. and van de Grift, P. 1996. History and science of knots. <i>River Edge</i> 15(3), 361-369.	An overview of the physics behind and the application for knots.					
Eaton, T. A. and Clark, A. J. 2015. Muscle articulation: flexible jaw joints made of soft tissue. <i>Integrative and Comparative Biology</i> 55, 1-12.	Survey of animals that use soft tissue to build jaw joints, including the morphology, behavior, physiology, and feeding behavior of hagfish.	https://academic.oup.com/iand/article/55/2/1/1248611/Music-Articulation-Flexible-Jaw-Joints-Made-of-Soft-Tissue				
Wells, R. M. C. and Forrester, M. E. 1989. Short communication: dependence of blood viscosity on haematocrit and shear rate in a primitive vertebrate. <i>J Exp Biol</i> 145, 483-487.	This paper investigates the biomechanics of blood circulation in hagfish.	http://jeb.biologists.org/content/145/3/483.full.pdf				

Clark, A. J., Marzillo, F. J., and Summers, A. P. 2010. A soft organ for a forceful bite: novel features of the feeding musculature in Atlantic hagfish, <i>Myxine glutinosa</i> . <i>Zool</i> 113, 259-266.	This paper is an examination of motor patterns and musculature involved in feeding without a rigid support system in the jaw.	http://dx.doi.org/10.1093/zool/113.2.259		
Clark, A. J. and Summers, A. P. 2007. Morphology and kinematics of feeding in hagfish: specific functional adaptations of jaws. <i>The Journal of Experimental Biology</i> 210, 3897-3909.	This paper examines feeding morphology and kinematics in two hagfish species, <i>Epiplatys aoteana</i> and <i>Myxine glutinosa</i> .	http://faculty.washington.edu/ajclark/Research/Research_PDFs/2007_hagfish_feeding_mechanics.pdf		
Cole, J. J. 1907. A monograph of the general morphology of the myxineid fishes, based on a study of <i>Myxine</i> . Part II: The anatomy of the muscles. <i>Transactions of the Royal Society of Edinburgh</i> 45, 543-757.	Cole breaks the myxineid fishes down by muscle groups and analyzes the anatomy and physiology of those muscle groups.			
Dawson, I. A. 1963. The oral cavity, the 'jaw' and the horny tooth of <i>Myxine glutinosa</i> . <i>Rev. Mycol.</i> 23:1-251.	Dawson focuses on the oral cavity and other features associated with feeding. More specifically, he looks at the histology and musculature of the oral cavity.			
Dawson, I. A. 1969. The keratinized tooth of <i>Myxine glutinosa</i> : a histological, histochemical, ultrastructural and experimental study. <i>Acta Zoologica</i> 40, 15-64.	Dawson details histological and histochemical studies of hagfish teeth.	http://library.ualberta.ca/2658/1/sign?ref=http://search.elsevier.com/locate/0169-4728(69)40:1-64::FT		
Genhullis, S. and Bursack, P. 2002. Architecture of the integument in lower teleosts: functional morphology and evolutionary implications. <i>J. Morphol.</i> 253, 290-309.	This paper provides a comparative description of the integument of ganoid and rayganoid fishes.	http://onlinelibrary.wiley.com/doi/10.1002/jmor.10072		
Genhullis, S. and Vogel, F. 2002. Spatial arrangement of white muscle fibers and connective tissue in fishes. <i>Comp Biochem & Physiol</i> 133, 1813-1827.	This paper describes the arrangement of white muscle fibers and the framework that it creates for research on fish swimming mechanics.	http://www.sciencepress.com.au/science/article/pii/S1895413302001861		
Glover, C. N., Baskling, C., and Wood, C. M. 2011. Characterization of L-alanine and glycine absorption across the gut of an ancient vertebrate. <i>J. Comp Physiol B</i> 141, 765-771.	In vivo techniques were used to examine gastrointestinal nutrient uptake in Pacific hagfish, <i>Epiplatys aoteana</i> .	https://www.zoology.ualberta.ca/~woodcm/wordpress/content/uploads/2016/07/Glover-et-al-2011-JCPB.pdf		
Glover, C. N., Baskling, C., and Wood, C. M. 2011. Adaptions to in vivo feeding: novel nutrient acquisition pathways in an ancient vertebrate. <i>Proc R Soc B</i> 1-6.	This paper studies the ability of Pacific hagfish to absorb amino acids and nutrients across the gill and skin surfaces.	http://rspb.royalsocietypublishing.org/content/278/1721/3096.abstract		
McDonald, B. G., Cavalli, V., Cabrett, L., and Milligan, C. L. 1991. Acid-base regulation in the Atlantic hagfish, <i>Myxine glutinosa</i> . <i>J. Exp Biol</i> 164, 201-215.	This paper describes net transfer of acid equivalents to the external environment in hagfish.	https://pubs.scientificdata.org/1052284/1991/0008/164M134815480.pdf		
Muller, J. 1914. Vergleichende Anatomie der Myxineiden, des cyclonemalen und anichthyoiden Ganoiden. <i>Abhandlungen der Königl. Preuss. Akademie der Wissenschaften in Berlin</i> 65-540.	This paper looks at ontological and myological structures and functions in myxineids.	http://www.biodidacticlibrary.org/items/9290/page/10/muller14p		
Schultz, A. G., Gaffney, S. C., Clifford, A. M., and Goss, G. 2014. Phosphate absorption across multiple epithelia in the Pacific hagfish, <i>Epiplatys aoteana</i> . <i>Am J Physiol Regul Integr Comp Physiol</i> 307, R464-R472.	This study investigates the mechanism of phosphate absorption in hagfish.	http://ajrph.physiology.org/content/307/5/R464.full		
Uyeno, T. and Kato, W. M. 2007. Electromyography of the basal musculature of cephalopods (<i>Sepia bimaculata</i>) in a state of the muscle activation in support and movement. <i>J. Exp Biol</i> 210, 118-128.	This study uses electromyography to test previous predictions that basal mass acts as the effector and the joint.	https://www.researchgate.net/profile/Willem_Kato/publication/16382111_Electromyography_of_the_basal_musculature_of_cephepod_bimaculata_a_test_of_the_hypothesis_of_the_muscle_activation_in_support_and_movement/links/56278684526427836.pdf		
Wells, R. M. G., Farrow, M. E., Davison, W., Taylor, H. H., Davis, P. S., and Satchell, C. R. 1986. Blood oxygen transport in the three swimming hagfish, <i>Epiplatys aoteana</i> . <i>J. Exp Biol</i> 121, 43-53.	This paper studies the physiology of oxygen transport in the blood of hagfish.	https://www.researchgate.net/profile/Priscilla_Davis/publication/1816134_Blood_oxygen_transport_in_the_three_swimming_hagfish_Epiplatys_aoteana_cerata/links/5953d26240047708000.pdf		

Aiz, J. W. 1952. Nasal breathing in fishes and the evolution of internal nares. <i>Quart Rev Bio</i> 27, 366-377.	This research looks at the origin and evolution of internal nares of aquatic and terrestrial vertebrates.	http://www.jstor.org/stable/pdf/2893731.pdf	
Auster, P. J. and Barber, K. 2003. Atlantic hagfish exploit prey capture by other taxa. <i>Journal of Fish Biology</i> 68, 618-623.	This paper describes observations of Atlantic Hagfish utilizing encounter competition by feeding on prey captured by other species.	https://www.researchgate.net/profile/Peter_Auster/publication/21379277_Atlantic_hagfish_exploit_preay_captured_by_other_taxa/links/0046533461941407000000.pdf	
Ayres, H. and Jackson, C. M. 1901. Morphology of the myxinoidei. <i>J Morphol</i> 17, 185-226.	This paper gives an anatomical overview of the skeleton and musculature of myxinoidei.		
Bardele, D. 1991. First fossil hagfish (Myxinoidea): a record from the Pennsylvanian of Illinois. <i>Science</i> 254, 701-703.	This paper discusses a new hagfish fossil and describes how it supports the conservative geological history of myxinoidei.	http://search.proquest.com/docview/213547228?accountid=14800	
Butler, A. R. 2000. Sensory system evolution at the origin of craniates. <i>Philosophical Transactions: Biological Sciences</i> 355, 1309-1313.	This paper looks at the steps in the evolution of sensory systems during the transition from non-craniate invertebrates to craniates.	http://www.jstor.org/library/valkosta.edu/2048/stable/pdf/3066668.pdf	
Catchpole, T. L., Frid, C. L. J., and Gray, T. S. 2006. Importance of discards from the English Nephrops norvegicus fishery in the North Sea to marine scavengers. <i>Mar Ecol Prog Ser</i> 343, 23-26.	This paper discusses the positive effect that fishery discards have on marine scavenger populations.	https://www.researchgate.net/publication/250218699_Importance_of_discards_from_the_English_Nephrops_norvegicus_fishery_in_the_North_Sea_to_marine_scavengers	
Cerny, R., Kartell, M., Saska-Spangler, T., Bronner-Frazer, M., Yu, F., and Medeiros, D. M. 2010. Evidence for the prepattern/cooption model of vertebrate jaw evolution. <i>PNAS</i> 107, 17262-17267.	The authors examined expression of 12 genes in pharyngeal patterning and determined that gnathostome pharyngeal patterning developed prior to the evolution of jaws.	http://www.jstor.org/library/valkosta.edu/2048/stable/pdf/20779951.pdf	
Ellis, J. E., Rowe, S., and Lotze, H. K. 2015. Expansion of hagfish fisheries in Atlantic Canada and worldwide. <i>Fish Res</i> 161, 24-33.	This article discusses the history of hagfish fisheries and the sustainability of modern fisheries.	https://www.researchgate.net/publication/26405162_Expansion_of_hagfish_fisheries_in_Atlantic_Canada_and_worldwide	
Forsy, P. and Janvier, P. 1994. Evolution of the early vertebrates. <i>American Scientist</i> 82, 554-565.	The authors discuss recent fossil discoveries and what they reveal about modern species and their relationship to ancestors during the time of jaw development.	http://www.jstor.org/library/valkosta.edu/2048/stable/pdf/29775327.pdf	
Furbringer, P. 1875. Untersuchungen zur vergleichenden anatomie der muskulatur des kopfes der cyclostomen. <i>Anatomie Zeitschrift für Naturwissenschaft</i> 9, 1-92.	This paper investigates the comparative anatomy of cyclostome musculature.	http://www.biodiversitylibrary.org/item/35028?page=7#mode=1up	
Gai, Z., Donghua, P. C., Zhu, M., Janvier, P., and Stappanoni, M. 2011. Fossil jawless fish from China foreshadows early jawed vertebrate anatomy. <i>Nature</i> 476, 324-327.	Synchrotron radiation X-ray tomography was used to describe the cranial anatomy of galeaspids, an ostracoderm group, in order to study jaw evolution.	http://dx.doi.org/10.1038/nature09484	
Goldstein, D. J. 1961. On the origin and morphology of myoepithelial cells of apocrine sweat glands. <i>J Evol Form</i> , 301-309.	This paper describes the origin of apocrine sweat glands.	http://www.sciencedirect.com/science/article/pii/S0022022X15486046	
Griffith, R. W. 1994. The life of the first vertebrate. <i>Bio Science</i> 44, 408-417.	Griffith provides a possible story line for the transition from invertebrate to vertebrate species and discuss the position of hagfish in this story.	http://www.jstor.org/stable/pdf/1312464.pdf	
Heimberg, A. M., Cowper-Salari, R., Sermon, M., Donghua, P. C. J., and Peterson, K. J. 2010. MicroRNAs reveal the interrelationships of hagfish, lampreys, and gnathostomes and the nature of the ancestral vertebrate. <i>PNAS</i> 107, 19379-19383.	The authors prove that Cyclostomata is monophyletic through shared microRNA families in lampreys and hagfish.	http://www.jstor.org/library/valkosta.edu/2048/stable/pdf/25748682.pdf	
Janvier, P. 2008. Early jawless vertebrates and cyclostome origins. <i>Zoological Society of Japan</i> 85, 1045-1056.	Janvier discusses the current organization of cyclostomes in the phylogenetic tree while addressing discrepancies in organization techniques and the lack of fossil intermediates between lampreys and hagfish.	http://www.biocore.org/doi/full/10.2108/zju.25.1045	
Jensen, D. 1966. The hagfish. <i>Sci Amer</i> 83-90.	This article is a broad overview of hagfish anatomy, physiology, and behavior.		
Jungersen, J. M., Lombolt, J. P., Weber, R. E., and Maltz, H. 1998. The biology of hagfishes. London (UK). <i>Chapman and Hall</i> 1, 1-578.	This book covers the evolution, anatomy, and physiology of the hagfish.		
Karatani, S. 2004. Evolution of the vertebrate jaw: comparative embryology and molecular developmental biology reveal the factors behind evolutionary novelty. <i>J Anat</i> 205, 335-347.	This article investigates the evolution of the jaw through embryology and molecular developmental biology.	http://onlinelibrary.wiley.com/doi/10.1111/j.1469-7524.2004.00345.x/full	
Karatani, S. and Ota, K. G. 2008. Primitive versus derived traits in the developmental program of the vertebrate head: views from cyclostome developmental studies. <i>J Exp Zool</i> 310, 294-314.	Karatani and others study the early evolution of the vertebrate skull.	http://onlinelibrary.wiley.com/library/valkosta.edu/2048/doi/10.1002/ajz.b.21190.pdf	
Karatani, S., Nishizawa, Y., Harigome, N., and Shigetani, Y. 2001. Embryology of the lamprey and evolution of the vertebrate jaw: insights from the molecular and developmental perspectives. <i>Phil Trans R Soc Lond</i> 356, 1615-1632.	This paper investigates the evolution of the jaw in lampreys.	http://www.jstor.org/library/valkosta.edu/2048/stable/pdf/3066662.pdf	
Lauber, A. 1938. Die visceralmuskulatur der acranier, cyclostomen und fische. <i>Handbuch der vergleichenden Anatomie der Wirbeltiere</i> , 468-479.	This paper studies the visceral, cranial musculature of cyclostomes.		
Mallat, J. 1997. Crossing a major morphological boundary: the origin of jaws in vertebrates. <i>Zoology</i> 100, 128-140.	This paper reviews the origin of jaws in vertebrates.	https://www.researchgate.net/publication/285533178_Crossing_a_major_morphological_boundary_The_origin_of_jaws_in Vertebrates	
McLeod, R. J. and Wing, S. R. 2007. Hagfish in the New Zealand fjords are supported by chemosymbiosis of forest carbon. <i>Ecology</i> 88, 809-816.	This paper analyzes the contribution of chemosymbiosis to the hagfish diet.	https://www.researchgate.net/profile/Rebecca_McLeod/publication/6298836_Hagfish_in_the_New_Zealand_fjords_are_supported_by_chemosymbiosis_of_forest_carbon/links/5634f1508ac34c9c4db62.pdf	
Miscornum, M. M. and Stewart, A. L. 2006. A new species of giant seven-gilled hagfish (Myxinoidea: Ipnopneustes) from New Zealand. <i>Copeia</i> 2, 225-229.	This paper describes a new species of hagfish caught in New Zealand.	http://dx.doi.org/10.1111/j.1365-3113.2006.03691.x	
Miyashita, T. 2012. Comparative analysis of the anatomy of the myxinoidea and the ancestry of early vertebrate lineages. <i>University of Alberta Dept Biol Sci</i> , 1-407.	This thesis presents a comparative analysis of hagfish anatomy and discusses the classification of hagfish.	https://era.library.ualberta.ca/files/p89807a?Myashita_Tetsuo_Summer2012.pdf	
Ohi, Y., Fujimoto, S., Ota, K. G., and Karatani, S. 2013. On the peculiar morphology and development of the hypoglossal, glossopharyngeal and vagus nerves and hypobranchial muscles in the hagfish. <i>Zoological Letters</i> 1, 1-15.	This paper analyzes the morphology and development of various nerves in the hagfish.	https://zoology.ualberta.ca/files/p89807a?Myashita_Tetsuo_Summer2012.pdf	
Ota, K. G., Fujimoto, S., Ohi, Y., and Karatani, S. 2011. Identification of vertebra-like elements and their possible differentiation from sclerotomes in the hagfish. <i>Nature Communications</i> 2, 1-6.	This paper identifies vertebra-like elements in the hagfish.	http://www.nature.com/articles/ncomms1355	
Shimeld, S. M. and Donghua, P. C. J. 2012. Evolutionary crossroads in developmental biology: cyclostomes (lamprey and hagfish). <i>Development</i> 139, 2091-2099.	This paper summarizes the development of cyclostomes, highlighting the key species studied and experimental methods available.	http://palaeo.gly.bris.ac.uk/donghua/PDFs/2012Shimeld_Donghua_2012.pdf	
Smith, C. R. and Bacon, A. R. 2003. Ecology of whale falls at the deep-sea floor. <i>Oceanography and Marine Biology</i> 41, 311-334.	This paper discusses the ecological impact of whale falls at the sea floor, including acting as a food source for hagfish.	https://www.oest.hawaii.edu/oceanography/faculty/csmith/Files/Smith%20and%20Bacon%202003.pdf	
Soukup, V., Horacek, I., and Cerny, R. 2013. Development and evolution of the vertebrate primary mouth. <i>J Anat</i> 222, 79-99.	Primary mouth formation is reviewed from a comparative point of view in order to reveal its morphogenetic diversity among vertebrate clades.	http://onlinelibrary.wiley.com/doi/10.1111/j.1469-7524.2012.01540.x/full	
Strahan, R. 1963. The behavior of myxinoidei. <i>Acta Zoologica</i> 44, 73-102.	This paper studies hagfish behavior in locomotion, burrowing, feeding, and light response, and compares behavior of hagfish and lampreys.		
Uyeno, T. A. and Kier, W. M. 2005. Functional morphology of the cephalopod buccal mass: a novel joint type. <i>J Morphol</i> 264, 211-222.	The arrangement of the musculature and connective tissues of the buccal mass of the colored cephalopods was examined using dissection and histology.	http://labs.bio.unc.edu/kier/pdf/Uyeno_Kier_2005.pdf	
Vogel, F. and Gemballa, S. 2000. Locomotory design of 'cyclostome' fishes: spatial arrangement and architecture of myosepta and lamellae. <i>Acta Zoologica</i> 81, 267-283.	This paper compares the myoseptal system of gnathostome locomotion to that of hagfish and lampreys.		
Wlga, C. D., Wainwright, P. C., and Motta, P. J. 2000. Evolution of jaw depression mechanisms in aquatic vertebrates: insights from Chondrichthyes. <i>Biological Journal of the Linnean Society</i> 71, 165-185.	This paper reviews key findings on the feeding mechanisms of elasmobranchs and present new ideas on the origin of jaw depression mechanisms.	http://www.eyes.acadavis.edu/wainwrightlab/Wlga_et_al_2000.pdf	
Werthington, J. 1905. Contribution to our knowledge of the myxinoidei. <i>American Naturalist</i> 39, 625-663.	This paper gives an early, general description of hagfish.	http://www.jstor.org/stable/2454811?seq=4#page_scan_tab_contents	
Zhikun, G. and Min, Z. 2012. The origin of the vertebrate jaw: intersection between developmental biology-based model and fossil evidence. <i>Chin Sci Bull</i> 57, 3819-3828.	This paper investigates the origin of the vertebrate jaw and sparks debate over galeaspids vs ostracoderms as the better proxy for reconstructing the pre-gnathostome condition of the rostral part of the braincase.	http://dx.doi.org/10.1007/s11434-012-5372-z?l=trac	
Ziermann, J. M., Miyashita, T., and Diego, R. 2014. Cephalic muscles of cyclostomes (hagfish and lampreys) and Chondrichthyes (sharks, rays, and holocephalus): comparative anatomy and early evolution of the vertebrate head muscles. <i>Zoological Journal of the Linnean Society</i> 172, 771-802.	This paper provides a comparative analysis of the anatomy and evolution of cyclostome, chondrichthyan, and osteichthyan cephalic musculature.	http://onlinelibrary.wiley.com/doi/10.1111/zju.12138.pdf?v=1&t=159w05&c=0186f97361b5ca05250a088ec4d4517980578f	
Zintzen, V., Roberts, C. D., Anderson, M. J., Stewart, A. L., Struthers, C. D., and Harvey, Eann, S. 2011. Hagfish predatory behavior and slime defense mechanism. <i>Scientific Reports</i> 1, 1-6.	This article provides evidence to show that hagfish act as active predators while also providing observations on the mechanism of slime defense strategies.	https://pdfs.semanticscholar.org/308f/8884f68cfa459b69948bae59c62b8d4c.pdf	

Biewener, A. A. 1992. Biomechanics structures and systems a practical approach. New York (NY). <i>Oxford University Press</i> 1 , 1-283.	Textbook covering different structures and systems used in biomechanics research.			
Chirikjian, G. S. and Burdick, J. W. 1994. A modal approach to hyper-redundant manipulator kinematics. <i>IEEE Transactions on Robotics and Automation</i> 10 , 343-354.	This paper presents efficient kinematic modeling techniques for "hyper-redundant" robots.	http://authors.library.caltech.edu/6494/1/CH11ecetra94.pdf		
Gregg, J. 2014. Method for sea floor drilling using hagfish slime as drilling fluid additive. <i>Gregg Drilling & Testing Inc.</i> US 8,912,131 B2.	This United States patent details the use of hagfish slime in a process for sea floor drilling.	https://www.google.com/patents/WO2013059031A1?cl=en		
Janvier, P. 1996. The dawn of the vertebrates: characters versus common ascent in the rise of current vertebrate phylogenies. <i>Paleontology</i> 39 , 259-287.	Techniques in creating phylogenetic trees are debated using the placement of ostracoderms and hagfish based on characters vs. common ascent.	http://cdn.palass.org/publications/palaeontology/volume_39/pdf/vol39_part2_pp259-287.pdf		
Vincent, J. F. V. 1992. Biomechanics materials a practical approach. New York (NY). <i>Oxford University Press</i> 1 , 1-247.	Textbook overview of materials and formulas commonly studied in the field of biomechanics.			